

References

Albanese, J.R., et al. 1984. Geological and Hydrogeologic Research at the Western New York Nuclear Service Center, West Valley, New York. Final Report, August 1982-December 1983. U.S. Nuclear Regulatory Commission Report, NUREG/CR-3782.

Bergeron, M. P., et al. 1987. Geohydrologic Conditions at the Nuclear Fuels Reprocessing Plant and Waste Management Facilities at the Western New York Nuclear Services Center, Cattaraugus County, New York. U.S. Geological Survey Water Resources Investigations Report 85-4145.

Bergeron, M.P., and E.F. Bugliosi. 1988. Groundwater Flow Near Two Radioactive Waste Disposal Areas at the Western New York Nuclear Service Center, Cattaraugus County, New York – Results of Flow Simulation. U.S. Geological Survey Water Resources Investigations Report 86-4351.

Broughton, J.G., et al. 1966. Geologic Map of New York State. New York State Museum and Science Service Map and Chart Series No.5.

Committee on Biological Effects of Ionizing Radiations. 1980. The Effects on Populations of Exposure to Low Levels of Ionizing Radiation. BEIR V. Washington: National Academy Press.

Comprehensive Environmental Response, Compensation, and Liability Act of 1980. December 11, 1980. Public Law 96-150, 94 Stat 2767, Title 26.

Dames & Moore. May 1991. Summary Report, RCRA Well Installation Program 1989-1990.

Dooley, D. A., and J. Prowse. 1990. Radiological Parameters for Assessment of WVDP Activities. Rev. 2. WVDP-065.

Dunning, Donald E. nd. Estimates of Internal Dose Equivalent from Inhalation and Ingestion of Selected Radionuclides. Revised. WIPP-DOE-176.

International Commission on Radiological Protection.

- Recommendations of the International Commission on Radiological Protection – Permissible Dose for Internal Radiation. 1959. ICRP Publication 2. Oxford: Pergamon Press.
- Report of the Task Group on Reference Man. 1975. ICRP Publication 23. Oxford: Pergamon Press.
- Recommendations of the International Commission on Radiological Protection. 1977. ICRP Publication 26. Oxford: Pergamon Press.
- Recommendations of the International Commission on Radiological Protection – Limits for Intakes of Radionuclides by Workers. 1979. ICRP Publication 30. Oxford: Pergamon Press.

Kalbeitzer, F. March 20, 1987. U.S. Department of Energy, Radiological and Environmental Sciences Laboratory, Idaho National Engineering Laboratory. Personal communication.

LaFleur, R.G. 1979. Glacial Geology and Stratigraphy of Western New York Nuclear Service Center and Vicinity, Cattaraugus and Erie Counties, New York. U.S. Geological Survey Open File Report 79-989.

Marchetti, S. December 17, 1982. Tritium in Groundwater. Letter (WD:82:0361) to W. Hannum, Department of Energy, West Valley Project Office.

Moore, R.E., et al. June 1979. AIRDOS-EPA: A Computerized Methodology for Estimating Environmental Concentrations and Dose to Man from Airborne Releases of Radionuclides. ORNL-5532.

References

National Council on Radiation Protection and Measurements.

- Environmental Radiation Measurements. December 1979. NCRP-50. Washington, D.C.
- Recommendations on Radiation Exposure Limits. July 1985. Draft report. Washington, D.C.
- Ionizing Radiation Exposure of the Population of the United States. 1987. NCRP-93. Bethesda, Maryland.

Oak Ridge National Laboratory. May 1980. User's Manual for LADTAP II - A Computer Program for Calculating Radiation Exposure to Man from Routine Release of Nuclear Reactor Liquid Effluents. NUREG/CR-1276.

Resource Conservation and Recovery Act of 1976. October 23, 1976. Public Law 94-580, 90 Stat. 2795, Title 42.

Rickard, L.V. 1975. Correlation of the Silurian and Devonian Rocks in New York State. New York State Museum and Science Service Map and Chart Series No. 24.

Sheppard, M.I., and D.H.Thibault. 1990. Default Soil Solid/Liquid Partition Coefficients, K_{ds} , for Four Major Soil Types: A Compendium. Health Physics. 59 (no. 4): 471-482.

Simpson, D.B., and B.L. McGill. 1980. LADTAP II: A Computer Program for Calculating Radiation Exposure to Man from Routine Release of Nuclear Reactor Liquid Effluents. Technical Data Management Center. ORNL/NUREG/TDMC-1.

Standish, P.N. 1985. Closure of the Construction Landfill Site. Letter (WD:85:0434) to W.H. Hannum, Department of Energy, West Valley Project Office.

Tesmer, I.H. 1975. Geology of Cattaraugus County, New York. Buffalo Society of Natural Sciences Bulletin, Vol. 27.

Tseng, J.C. November 4, 1975. Clarification of Applicable Radiation Protection Standards for the Public and the Environment. Memorandum from EH-231, Office of Environmental Guidance and Compliance, Washington, D.C.

Superfund Amendments and Reauthorization Act of 1986. October 17, 1986. Public Law 99-499, 100 Stat. 1613, Title 10.

U.S. Department of Energy.

- A Guide for Environmental Radiological Surveillance at U.S. Department of Energy Installations. July 1981. DOE/EP-0023. Washington, D.C.
- A Guide for Effluent Radiological Measurements at DOE Installations. July 1983. DOE/EP-0096. Washington, D.C.
- Environmental Safety and Health Program for Department of Energy Operations. DOE Order 5480.1B. September 1986. Washington, D.C.
- Environmental Protection, Safety and Health Protection Information Reporting Requirements. 1981. DOE Order 5484.1. Washington, D.C.
- General Environmental Protection Program. DOE Order 5400.1. November 1988. Washington, D.C.
- Radiation Protection of the Public and Environment. 1990. DOE Order 5400.5. Washington, D.C.
- Radiological Effluent Monitoring and Environmental Surveillance. 1990. Draft DOE Order 5400.6. Washington, D.C.

References

U. S. Environmental Protection Agency.

- National Interim Primary Drinking Water Regulations. 1976. EPA-570/9-76-003. Washington, D.C. : Office of Water Supply.
- National Emission Standard for Hazardous Air Pollutants; Standards for Radionuclides. 40 CFR 61 1983. Washington, D.C.: U.S. Government Printing Office.
- Drinking Water Guidelines. 40 CFR 141, National Secondary Drinking Water Regulations, Subpart B, *Maximum Contaminant Levels*, July 1, 1984.
- Drinking Water Guidelines. 40 CFR 143, National Secondary Drinking Water Regulations, Section 143.3, *Secondary Maximum Contaminant Levels* (1984b).
- Groundwater Monitoring Technical Enforcement Guidance Document. 1986. OWSER-9950.1. Washington, D.C.
- Statistical Analysis of Groundwater Monitoring Data at RCRA Facilities. 1989. Interim Final Guidance. EPA/530-SW-89-026. Washington, D.C.

U.S. Nuclear Regulatory Commission.

- Regulatory Guide 1.109: Calculation of Annual Doses to Man from Routine Releases of Reactor Effluents for the Purpose of Evaluating Compliance with 10 CFR Part 50, Appendix I. October 1977.
- Regulatory Guide 1.111: Methods for Estimating Atmospheric Transport and Dispersion of Gaseous Effluents in Routine Releases from Light-Water-Cooled Reactors. July 1977.
- Standard for Protection Against Radiation. Proposed rule, 10 CFR Parts 19, 20, et al., Federal Register, 51, No. 6. February 9, 1986.

West Valley Demonstration Project. 1986. WVDP Radioactive Air Emissions Permit Application General Information. Submitted to EPA Region 2.

West Valley Nuclear Services, Inc.

- West Valley Demonstration Project Safety Analysis Report. Vol. 1 Supplements. June 1985.
- 1986 Environmental Monitoring Report, West Valley Demonstration Project. March 1987. WVDP-040.
- 1987 Effluent and On-Site Discharge Report, West Valley Demonstration Project. March 1988.
- 1987 Environmental Monitoring Report, West Valley Demonstration Project. March 1988. WVDP-040.
- West Valley Demonstration Project Site Environmental Monitoring Report for Calendar Year 1988. May 1989.
- West Valley Demonstration Project Site Environmental Report for Calendar Year 1989. May 1990.
- Groundwater Protection Management Program. April 1990. WVDP-091.

Yager, R.M. 1987. Simulation of Groundwater Flow near the Nuclear Fuel Reprocessing Facility at the Western New York Nuclear Service Center, Cattaraugus County, New York. 1987. 85-4308. Ithaca, New York: U.S. Geological Survey.

Glossary

ALLUVIUM. Sedimentary material deposited by flowing water such as a river.

ALLUVIAL FAN. A cone-shaped deposit of alluvium made by a stream where it runs out onto a level plain.

AQUIFER. A water-bearing unit of permeable rock or soil that will yield water in usable quantities to wells. *Confined aquifers* are bounded above and below by less permeable layers. Groundwater in a confined aquifer is under a pressure greater than the atmospheric pressure. *Unconfined aquifers* are bounded below by less permeable material, but are not bounded above. The pressure on the groundwater in an unconfined aquifer at the top of the aquifer is equal to that of the atmosphere.

AQUITARD. A relatively impervious and semiconfining geologic formation which, compared to an aquifer, transmits water at a very slow rate .

BACKGROUND RADIATION. Includes both natural and manmade radiation such as cosmic radiation and radiation from naturally radioactive elements and from commercial sources and medical procedures.

BECQUEREL (BQ). A unit of radioactivity equal to one nuclear transformation per second.

CLASS A, B, AND C LOW-LEVEL WASTE. Waste classifications from the Nuclear Regulatory Commission's 10 CFR Part 61 rule. Maximum concentration limits are set for specific isotopes. Class A waste disposal is minimally restricted with respect to the form of the waste. Class B waste must meet more rigorous requirements to ensure physical stability after disposal. Greater concentration limits are set for the same isotopes in Class C waste, which also must meet physical stability requirements. Moreover, special measures must be taken at the disposal facility to protect against inadvertent intrusion.

CONFIDENCE COEFFICIENT OR FACTOR. The chance or probability, usually expressed as a percentage, that a confidence interval includes some defined parameter of a population. The confidence coefficients usually associated with confidence intervals are 90%, 95%, and 99%.

COSMIC RADIATION. High-energy subatomic particles from outer space that bombard the earth's atmosphere. Cosmic radiation is part of natural background radiation.

COUNTING ERROR. The variability caused by the inherent random nature of radioactive disintegration and the detection process.

CURIE (Ci). A unit of radioactivity equal to 37 billion (3.7×10^{10}) nuclear transformations per second.

DETECTION LEVEL. The minimum concentration of a substance that can be measured with a 99% confidence that the analytical concentration is greater than zero.

DERIVED CONCENTRATION GUIDE (DCG). Concentrations of radionuclides in air and water in which a person continuously exposed and inhaling 8400 m³ of air or ingesting 730 liters of water per year would receive an annual effective dose equivalent of 100 mrem per year from either mode of

Glossary

exposure. The committed dose equivalent is included in the DCGs for radionuclides with long half-lives (see appendix B).

DISPERSION. The process whereby solutes are spread or mixed as they are transported by groundwater as it moves through sediments.

DOSIMETER. A portable device for measuring the total accumulated exposure to ionizing radiation.

DOWNGRADIENT. The direction of water flow from a reference point to a selected point of interest (see GRADIENT).

EFFECTIVE DOSE. See EFFECTIVE DOSE EQUIVALENT under RADIATION DOSE.

EFFLUENT. Flowing out or forth; an outflow of waste. In this report, effluent refers to the liquid or gaseous waste streams released into the environment from the facility.

EFFLUENT MONITORING. Sampling or measuring specific liquid or gaseous effluent streams for the presence of pollutants.

EXPOSURE. Subjecting a target (usually living tissue) to radiation.

FALLOUT. Radioactive materials mixed into the earth's atmosphere. Fallout constantly precipitates onto the earth.

GRADIENT. Change in value of one variable with respect to another variable, especially vertical or horizontal distance.

GROUNDWATER. Subsurface water in the pore spaces of soil and geologic units.

HALF-LIFE. The time in which half the atoms of a radionuclide disintegrate into another nuclear form. The half-life may vary from a fraction of a second to thousands of years.

HIGH-LEVEL WASTE (HLW). The highly radioactive waste material that results from the reprocessing of spent nuclear fuel, including liquid waste produced directly in reprocessing and any solid waste derived from the liquid, that contains a combination of transuranic waste and fission products in concentrations sufficient to require permanent isolation.

HYDRAULIC CONDUCTIVITY. The ratio of flow velocity to driving force for viscous flow under saturated conditions of a specified liquid in a porous medium; the ratio describing the rate at which water can move through a permeable medium.

ION. An atom or group of atoms with an electric charge.

ION EXCHANGE. The reversible exchange of ions contained in solution with other ions that are part of the ion-exchange material.

ISOTOPE. Different forms of the same chemical element that are distinguished by having different numbers of neutrons in the nucleus. An element can have many isotopes. For example, the three isotopes of hydrogen are protium, deuterium, and tritium.

Glossary

KAME DELTA. A conical hill or short irregular ridge of gravel or sand deposited in contact with glacier ice.

LACUSTRINE SEDIMENTS. A sedimentary deposit consisting of material pertaining to, produced by, or formed in a lake or lakes.

LEACHED HULLS. Stainless steel cladding that remains after acid dissolution of spent fuel.

LOW-LEVEL WASTE. Radioactive waste not classified as high-level waste, transuranic waste, spent fuel, or uranium mill tailings (see CLASS A,B,C LOW-LEVEL WASTE).

MAXIMALLY EXPOSED INDIVIDUAL. A hypothetical person who remains in an uncontrolled area who would, when all potential routes of exposure from a facility's operations are considered, receive the greatest possible dose equivalent.

MEAN. The average value of a series of measurements.

MILLIREM (MREM). A unit of radiation dose equivalent that is equal to one one-thousandth of a rem. An individual member of the public can receive up to 500 millirems per year according to DOE standards. This limit does not include radiation received for medical treatment or the 100 to 360 mrem that people receive annually from background radiation.

MINIMUM DETECTABLE CONCENTRATION. The smallest amount or concentration of a radioactive or nonradioactive element that can be reliably detected in a sample.

MIXED WASTE. A waste that is both radioactive and hazardous. Also referred to as RADIOACTIVE MIXED WASTE (RMW).

OUTFALL. The end of a drain or pipe that carries waste water or other effluents into a ditch, pond, or river.

PARTICULATES. Solid particles and liquid droplets small enough to become airborne.

PERSON-REM. The sum of the individual radiation dose equivalents received by members of a certain group or population. It may be calculated by multiplying the average dose per person by the number of persons exposed. For example, a thousand people each exposed to one millirem would have a collective dose of one person-rem.

PLUME. The distribution of a pollutant in air or water after being released from a source.

PROGLACIAL LAKE. A lake occupying a basin in front of a glacier; generally in direct contact with the ice.

RAD. Radiation absorbed dose. One hundred ergs of energy absorbed per gram.

RADIATION. The process of emitting energy in the form of rays or particles that are thrown off by disintegrating atoms. The rays or particles emitted may consist of alpha, beta, or gamma radiation.

Glossary

- **ALPHA RADIATION.** The least penetrating type of radiation. Alpha radiation can be stopped by a sheet of paper or outer dead layer of skin.
- **BETA RADIATION.** Electron emitted from a nucleus during fission and nuclear decay. Beta radiation can be stopped by an inch of wood or a thin sheet of aluminum.
- **GAMMA RADIATION.** A form of electromagnetic, high-energy radiation emitted from a nucleus. Gamma rays are essentially the same as x-rays and require heavy shielding such as lead, concrete, or steel to be stopped.
- **INTERNAL RADIATION.** Radiation originating from a source within the body as a result of the inhalation, ingestion, or implantation of natural or manmade radionuclides in body tissues.

RADIATION DOSE.

- **ABSORBED DOSE.** The amount of energy deposited by radiation in a given amount of material. Absorbed dose is measured in rads.
- **COLLECTIVE DOSE EQUIVALENT.** The sum of the dose equivalents for individuals comprising a defined population. The per capita dose equivalent is the quotient of the collective dose equivalent divided by the population (see PERSON-REM).
- **COMMITTED DOSE EQUIVALENT (CDE).** The effective dose equivalent from an intake of radionuclides delivered over a period of 50 years following the intake.
- **CUMULATIVE DOSE EQUIVALENT.** The total dose one could receive in a period of fifty years following release of radionuclides to the environment, including the dose that could occur as a result of residual radionuclides remaining in the environment beyond the year of release.
- **DOSE EQUIVALENT.** The product of the absorbed dose, the quality factor, and any other modifying factors. The dose equivalent is a quantity for comparing the biological effectiveness of different kinds of radiation on a common scale. The unit of dose equivalent is the rem.
- **EFFECTIVE DOSE EQUIVALENT.** The sum over all organs of dose equivalents (from internal and external radiation) to each organ, multiplied by the appropriate weighting factor for that organ.

RADIOACTIVITY. A property possessed by some elements such as uranium whereby alpha, beta, or gamma rays are spontaneously emitted.

RADIOISOTOPE. A radioactive isotope of a specified element. Carbon-14 is a radioisotope of carbon. Tritium is a radioisotope of hydrogen.

Glossary

RADIONUCLIDE. A radioactive nuclide. Radionuclides are variations (isotopes) of elements. They have the same number of protons and electrons but different numbers of neutrons, resulting in

different atomic masses. There are several hundred known nuclides, both manmade and naturally occurring.

REM. An acronym for Roentgen Equivalent Man. A unit of radiation exposure that indicates the potential effect on human cells.

SIEVERT. A unit of dose equivalent from the International System of Units. Equal to one joule per kilogram.

SPENT FUEL. Nuclear fuel that has been exposed in a nuclear reactor; this fuel contains uranium, activation products, fission products, and plutonium.

STANDARD DEVIATION. An indication of the dispersion of a set of results around their average.

THERMOLUMINESCENT DOSIMETER (TLD). A device that luminesces upon heating after being exposed to radiation. The amount of light emitted is proportional to the amount of radiation to which the luminescent material has been exposed.

UPGRADIENT. Referring to the flow of water or air, it is analogous to upstream. A point that is "before" an area of study that is used as a baseline for comparison with downstream data. See **GRADIENT** and **DOWNGRADIENT**.

WATERSHED. The area contained within a drainage divide above a specified point on a stream.

WATER TABLE. The upper surface in a body of groundwater. The surface in an unconfined aquifer or confining bed at which the pore water pressure is equal to atmospheric pressure.

WHOLE-BODY DOSE. A radiation dose that involves exposure of the entire body.

Acronyms

ANOVA. Analysis of Variance

ALARA. As Low As Reasonably Achievable

BEIR. Committee on Biological Effects of Ionizing Radiation

CDDL. Construction and Demolition Debris Landfill (formerly the “cold dump”)

CERCLA. Comprehensive Environmental Response, Compensation, and Liability Act

CSS. Cement Solidification System

DCG. Derived Concentration Guide

DE. Dose Equivalent

DOE. Department of Energy

DOE-HQ. Department of Energy, Headquarters Office

DOE-ID. Department of Energy, Idaho Operations

EA. Environmental Assessment

EDE. Effective Dose Equivalent

EE. Environmental Evaluation

EIS. Environmental Impact Statement

ELAP. Environmental Laboratory Accreditation Program

EML. Environmental Measurements Laboratory

EMSL. Environmental Monitoring Systems Laboratory (Las Vegas)

EPA. Environmental Protection Agency

FONSI. Finding of No Significant Impact

FSFCA. Federal and State Facilities Compliance Agreement

FY. Fiscal Year

HLW. High-level Radioactive Waste

ICRP. International Commission on Radiological Protection

INEL. Idaho National Engineering Laboratory

Acronyms

IRTS. Integrated Radwaste Treatment System

LLD. Lower Limit of Detection

LLW. Low-level Radioactive Waste

LLWTF. Low-level Liquid Waste Treatment Facility

LPS. Liquid Pre-treatment System

LWTS. Liquid Waste Treatment System

MDC. Minimum Detectable Concentration

NCRP. National Council on Radiation Protection and Measurements

NDA. Nuclear Regulatory Commission - licensed Disposal Area

NEPA. National Environmental Policy Act

NESHAP. National Emission Standards for Hazardous Air Pollutants

NIST. National Institute of Standards and Technology

NFS. Nuclear Fuel Services Company, Inc.

NOI. Notice of Intent

NRC. Nuclear Regulatory Commission

NWPA. Nuclear Waste Policy Act

NYSDEC. New York State Department of Environmental Conservation

NYSDOH. New York State Department of Health

NYSERDA. New York State Energy Research and Development Authority

NYSGS. New York State Geological Survey

OSR. Operational Safety Requirement

QA. Quality Assurance

QAP. Quality Assurance Program

QC. Quality Control

RCRA. Resource Conservation and Recovery Act

Acronyms

RMW. Radioactive Mixed Waste

SAR. Safety Analysis Report

SARA. Superfund Amendments and Reauthorization Act

SDA. (New York) State-licensed Disposal Area

SI. International System of Units

SPDES. State Pollutant Discharge Elimination System

STS. Supernatant Treatment System

SWMU. Solid Waste Management Unit

SSWMU. Super Solid Waste Management Unit

TLD. Thermoluminescent Dosimeter

USGS. U.S. Geological Survey

WNYNSC. Western New York Nuclear Service Center

WVDP. West Valley Demonstration Project

WVNS. West Valley Nuclear Services Co., Inc.

WVPO. West Valley (DOE) Project Office

Abbreviations for Units of Measure

	<u>Symbol</u>	<u>Name</u>		<u>Symbol</u>	<u>Name</u>
Radioactivity	Ci	curie	Volume	cm ³	cubic centimeter
	mCi	millicurie(1E-03Ci)		L	liter
	μCi	microcurie(1E-06Ci)		mL	milliliter
	nCi	nanocurie (1E-09 Ci)		m ³	cubic meter
	pCi	picocurie (1E-12 Ci)		ppm	parts per million
	fCi	femtocurie (1E-15Ci)		ppb	parts ber billion
	aCi	attocurie (1E-18 Ci)			
Bq	becquerel (27 pCi)				
	<u>Symbol</u>	<u>Name</u>		<u>Symbol</u>	<u>Name</u>
Dose	Sv	sievert (100 rems)	Time	y	year
	Gy	gray (100 rads)		d	day
				h	hour
		m		minute	
				s	second
	<u>Symbol</u>	<u>Name</u>		<u>Symbol</u>	<u>Name</u>
Length	m	meter	Area	ha	hectare (10,000 m ²)
	km	kilometer (1E + 03)			
	cm	centimeter (1E-02 m)			
	mm	millimeter (1E-03 m)			
	μm	micrometer (1E-06 m)			
	<u>Symbol</u>	<u>Name</u>			
Mass	g	gram			
	kg	kilogram (1E + 03 g)			
	mg	milligram (1E-03)			
	μg	microgram (1E-06 g)			
	ng	nanogram (1E-09 g)			
t	metric ton (10 ³ kg)				

Distribution

R. Natoli	DOE-HQ	T. DeBoer	NYSERDA
T. McIntosh	DOE-HQ	S. Harbison	NYSERDA
H. Walter	DOE-HQ	T. Sonntag	NYSERDA
J. Barry	DOE-ID	R. Fakudiny	NYSGS
B. Bowhan	DOE-ID	F. Galpin	USEPA-Washington, D.C.
M.B. Hinman	DOE-ID	F. Caporcuscio	USEPA-Region 2
D.B. Engelman	DOE-ID	P. Giardina	USEPA-Region 2
T. Burns	DOE-ID	J. Gorman	USEPA-Region 2
T. Rowland	DOE-WVPO	R. Novitski	USGS
D. Hurt	NRC-HQ	A. Stevens	SNIHD
M. Austin	NRC-Region 1	C. Halgas	CCHD
J. Roth	NRC-Region 1	W. Paxon	U.S. Congressman, 31st Dist.
P. Counterman	NYSDEC-Albany	A. Houghton	U.S. Congressman, 34th Dist
P. Merges	NYSDEC-Albany	D. Moynihan	U.S. Senator, New York
R. Baker	NYSDEC-Region 9	A. D'Amato	U.S. Senator, New York
B. Bartz	NYSDEC-Region 9	J. Present	New York Senator, 56th Dist.
P. Eisman	NYSDEC-Region 9	P. McGee	New York Assemblyman, 149th Dist.
M. Jackson	NYSDEC-Region 9		
B. Ignatz	NYSDOH-Buffalo	Concord Public Library	
K. Rimawi	NYSDOH-Albany	Springville, New York	

*News release summary

Community Relations, WVNS (Technical File)

Buffalo News, Buffalo, New York *

Salamanca Republican Press, Salamanca, New York *

Springville Journal, Springville, New York *