

**1995 Monitoring Program
Environmental Surveillance:**

On-site Groundwater

Sample Location Code	Monitoring/Reporting Requirements	Sampling Type/Medium	Collection Frequency	Total Annual Sample Collections	Analyses Performed/ Composite Frequency
Maintenance Shop Leach Fields (SSWMU #5)	Groundwater monitoring wells around site super solid waste management units (SSWMUs)* <u>Reported in:</u> • SER • Quarterly Groundwater Reports	Grab liquid	See Tables 3-1, 3-2, and Appendix E	See Tables 3-1, 3-2, and Appendix E	See Tables 3-1, 3-2, and 3-3
WNW 0501 U 0502		Direct field measurement of sample discharge water			
Low-level Waste Storage Area (SSWMU #6)					
WNW 0601 0602 0603 U 0604 0605 8607 U 8608 U 8609 U					
Chemical Process Cell Waste Storage Area (SSWMU #7)					
WNW 0701 U 0702 C 0703 0704 0705 C 0706 U 0707					

NOTE: "U" designates upgradient, "B" designates background, and "C" designates crossgradient wells; the remainder are downgradient.

* The groundwater monitoring program was revised in May 1995 after a review of results from previous years of sampling. The program in place at the end of 1995 is presented in the "Groundwater Monitoring Plan" (WVDP-239).

Sampling Rationale

On-site Groundwater	<p>DOE Order 5400.1, IV.9; DOE/EH-0173T, 5.10.1.3; RCRA 3008(h) Order on Consent.</p> <p>The on-site WVDP groundwater monitoring program focuses on radiological and chemical surveillance of both active and inactive super solid waste management units (SSWMUs). The program allows for the determination of water quality. In addition, using wells situated hydraulically upgradient (background) and downgradient of SSWMUs allows for both detection of groundwater contamination and evaluation of the effects associated with the individual SSWMUs.</p> <p>Groundwater protection is addressed in WVDP-091, "Groundwater Protection Management Program." Groundwater monitoring as detailed in WVDP-239, "Groundwater Monitoring Plan," is applicable to the 1996 program.</p>
SSWMU #5	<p>Maintenance shop sanitary leach field, formerly used by NFS and WVNS to process domestic sewage generated by the maintenance shop.</p>
SSWMU #6	<p>The low-level waste storage area includes metal and fabric structures housing low-level radioactive wastes being stored for future disposal.</p>
SSWMU #7	<p>The chemical process cell (CPC) waste storage area contains packages of pipes, vessels, and debris from decontamination and cleanup of the chemical process cell in the former reprocessing plant.</p>

**1995 Monitoring Program
Environmental Surveillance:**

On-site Groundwater

Sample Location Code	Monitoring/Reporting Requirements	Sampling Type/Medium	Collection Frequency	Total Annual Sample Collections	Analyses Performed/Composite Frequency
Construction and Demolition Debris Landfill (CDDL) (SSWMU #8)	Groundwater monitoring wells around site super solid waste management units (SSWMUs)* <u>Reported in:</u> • SER • Quarterly Groundwater Reports	Grab liquid	See Tables 3-1, 3-2, and Appendix E	See Tables 3-1, 3-2, and Appendix E	See Tables 3-1, 3-2, and 3-3
WNW 0801 U 0802 0803 0804 8603 U 8612 WNGSEEP WNDMPNE		Direct field measurement of sample discharge water			
NRC-licensed Disposal Area (NDA) (SSWMU #9)					
WNW 0901 U 0902 U 0903 0904 0905 0906 0907 0908 U 0909 0910 8610 8611 WNNDATR					
IRTS Drum Cell (SSWMU #10)					
WNW 1001 U 1002 1003 1004 1005 U 1006 1007 1008b B 1008c B					

NOTE: "U" designates upgradient, "B" designates background, and "C" designates crossgradient wells; the remainder are downgradient.

* The groundwater monitoring program was revised in May 1995 after a review of results from previous years of sampling. The program in place at the end of 1995 is presented in the "Groundwater Monitoring Plan" (WVDP-239).

Sampling Rationale

On-site Groundwater	<p>DOE Order 5400.1, IV.9; DOE/EH-0173T, 5.10.1.3; RCRA 3008(h) Order on Consent.</p> <p>The on-site WVDP groundwater monitoring program focuses on radiological and chemical surveillance of both active and inactive super solid waste management units (SSWMUs). The program allows for the determination of water quality. In addition, using wells situated hydraulically upgradient (background) and downgradient of SSWMUs allows for both detection of groundwater contamination and evaluation of the effects associated with the individual SSWMUs.</p> <p>Groundwater protection is addressed in WVDP-091, "Groundwater Protection Management Program." Groundwater monitoring as detailed in WVDP-239, "Groundwater Monitoring Plan," is applicable to the 1996 program.</p>
SSWMU #8	<p>Construction and demolition debris landfill (CDDL), used by NFS and the WVDP to dispose of nonhazardous and nonradioactive materials.</p>
SSWMU #9	<p>The NRC-licensed disposal area (NDA) contains radioactive wastes generated by NFS and the WVDP.</p>
SSWMU #10	<p>The integrated radioactive waste system (IRTS) treatment drum cell stores cement-stabilized low-level radioactive waste.</p>

**1995 Monitoring Program
Environmental Surveillance:**

On-site Groundwater

Sample Location Code	Monitoring/Reporting Requirements	Sampling Type/Medium	Collection Frequency	Total Annual Sample Collections	Analyses Performed/Composite Frequency
State-licensed Disposal Area (SSWMU #11) WNW 1101a U 1101b U 1101c U 1102a 1102b 1103a 1103b 1103c 1104a 1104b 1104c 1105a 1105b 1106a U 1106b U 1107a 1108a U 1109a U 1109b U 1110a 1111a	Groundwater monitoring points around site super solid waste management units (SSWMUs) <u>Reported in:</u> • SER	Grab liquid	Per NYSERDA*	Per NYSERDA*	Per NYSERDA*
Fuel Storage Area (Not a SSWMU) WNW 8613A C 8613B C 8613C	<u>Reported in:</u> • SER • Quarterly Groundwater Reports	Grab liquid Direct field measurement of sample discharge water	See Tables 3-1, 3-2, and Appendix E	See Tables 3-1, 3-2, and Appendix E	See Tables 3-1, 3-2, and 3-3
Well Points (Not in a SSWMU) WP-A WP-C WP-D WP-E WP-F WP-G WP-H	Well points downgradient of main plant <u>Reported in:</u> • SER	Grab liquid	See Tables 3-1, 3-2, and Appendix E	See Tables 3-1, 3-2, and Appendix E	See Tables 3-1, 3-2, and 3-3

NOTE: "U" designates upgradient, "B" designates background, and "C" designates crossgradient wells; the remainder are downgradient.

* SDA wells are sampled by NYSERDA; therefore, frequencies and analyses are not included in this summary of the WVDP program. Data are presented in Appendix F.

Sampling Rationale

On-site Groundwater	<p>DOE Order 5400.1, IV.9; DOE/EH-0173T, 5.10.1.3; RCRA 3008(h) Order on Consent.</p> <p>The on-site WVDP groundwater monitoring program focuses on radiological and chemical surveillance of both active and inactive super solid waste management units (SSWMUs). The program allows for the determination of water quality. In addition, using wells situated hydraulically upgradient (background) and downgradient of SSWMUs allows for both detection of groundwater contamination and evaluation of the effects associated with the individual SSWMUs.</p> <p>Groundwater protection is addressed in WVDP-091, "Groundwater Protection Management Program." Groundwater monitoring as detailed in WVDP-239, "Groundwater Monitoring Plan," is applicable to the 1996 program.</p>
SSWMU #11	<p>The state-licensed disposal area (SDA) was operated by NFS as a commercial low-level disposal facility; it also received wastes from NFS reprocessing operations.</p>
Fuel Storage Area	<p>Monitors groundwater in the vicinity of the underground fuel storage tanks; this is not included in any of the SSWMUs.</p>
Well Points	<p>Monitor groundwater of known contamination in the north plateau area. All well points are downgradient of the main plant.</p>

**1995 Monitoring Program
Environmental Surveillance:**

Off-site Surface Water

Sample Location Code	Monitoring/Reporting Requirements	Sampling Type/Medium	Collection Frequency	Total Annual Sample Collections	Analyses Performed/Composite Frequency
WFBCTCB Buttermilk Creek, upstream of Cattaraugus Creek confluence at Thomas Corners Road	Restricted surface waters receiving plant effluents <u>Reported in:</u> • MTAR • QEMDR • SER	Timed continuous composite liquid	→ Weekly	→ 52	→ pH, conductivity
				Weekly samples composited to 12	→ Monthly composite for gross alpha/beta, H-3
				Weekly samples composited to 4	→ Quarterly composite for gamma isotopic and Sr-90
WFFELBR Cattaraugus Creek at Felton Bridge	Unrestricted surface waters receiving plant effluents <u>Reported in:</u> • MTAR • QEMDR • SER	Timed continuous composite liquid	→ Weekly	→ 52	→ Gross alpha/beta, H-3, pH
				Weekly samples composited to 12	→ Flow-weighted monthly composite for gamma isotopic and Sr-90, gross alpha/beta, H-3
WFBCBKG Buttermilk Creek near Fox Valley (background)	Unrestricted surface water background <u>Reported in:</u> • MTAR • QEMDR • SER	Timed continuous composite liquid	→ Weekly	→ 52	→ pH, conductivity
				Weekly samples composited to 12	→ Monthly composite for gross alpha/beta, H-3
			Weekly samples composited to 4	→ Quarterly composite for gamma isotopic, Sr-90, C-14, I-129, Pu/U isotopic, total U, Am-241, Tc-99	
		Grab liquid	→ Semiannual	→ 2	→ NPOC, TOX, Ca, Mg, Na, K, Ba, Mn, Fe, Cl, SO ₄ , NO ₃ -NO ₂ -N, F, HCO ₃ , CO ₃
WFBIGBR Cattaraugus Creek at Bigelow Bridge (background)	Unrestricted surface water background <u>Reported in:</u> • MTAR • QEMDR • SER	Grab liquid	→ Monthly	→ 12	→ Gross alpha/beta, H-3, Sr-90, and gamma isotopic

Monthly composite at **WFBCTCB**, **WFBCBKG**, and **WFFELBR** is also sent to NYSDOH.

Sampling Rationale

WFBCTCB DOE/EH-0173T, 5.10.1.1.

Buttermilk Creek is the surface water receiving all WVDP effluents. **WFBCTCB** monitors the potential influence of WVDP drainage into Buttermilk Creek upstream of confluence with Cattaraugus Creek.

WFFELBR DOE/EH-0173T, 5.10.1.1.

Because Buttermilk Creek is the surface water that receives all WVDP effluents and empties into Cattaraugus Creek, **WFFELBR** monitors the potential influence of WVDP drainage into Cattaraugus Creek directly downstream of the confluence with Buttermilk Creek.

WFBCBKG DOE/EH-0173T, 5.10.1.1.

Monitors background conditions of Buttermilk Creek upstream of the WVDP. Allows comparison to downstream conditions.

WFBIGBR DOE/EH-0173T, 5.10.1.1.

Monitors background conditions of Cattaraugus Creek at Bigelow Bridge, upstream of the WVDP. Allows comparison to downstream conditions.

**1995 Monitoring Program
Environmental Surveillance:**

Off-site Drinking Water

Sample Location Code	Monitoring/Reporting Requirements	Sampling Type/Medium	Collection Frequency	Total Annual Sample Collections	Analyses Performed/Composite Frequency
<p>WFWEL series wells near the WVDP outside the WNYNSC perimeter</p> <p>WFWEL01 3.0 km WNW</p> <p>WFWEL02 1.5 km NW</p> <p>WFWEL03 4.0 km NW</p> <p>WFWEL04 3.0 km NW</p> <p>WFWEL05 2.5 km SW</p> <p>WFWEL06 (background) 29 km S</p> <p>WFWEL07 4.0 km NNE</p> <p>WFWEL08 2.5 km ENE</p> <p>WFWEL09 3.0 km SE</p> <p>WFWEL10 7.0 km N</p>	<p>Drinking water supply; → groundwater near facility*</p> <p><u>Reported in:</u></p> <ul style="list-style-type: none"> • MTAR • QEMDR • SER 	→ Grab liquid	→ Annual	→ 1 each location	→ Gross alpha/beta, H-3, gamma isotopic, pH, conductivity

* Off-site drinking water wells are not affected by the potential migration of contaminants in the subsurface at the WVDP.

Sampling Rationale

Off-site Drinking Water WFWEL Series	DOE 5400.1, IV.9; DOE/EH-0173T, 5.10.1.2. Eight of the ten listed off-site private residential drinking water wells represent the nearest unrestricted uses of groundwater close to the WVDP. The ninth sample (WFWEL10) is from a public water supply from deep wells. The tenth drinking water well, WFWEL06 , is located 29 kilometers south of the Project and is considered a background drinking water source.
---	---

**1995 Monitoring Program
Environmental Surveillance:**

Off-site Air

Sample Location Code	Monitoring/Reporting Requirements	Sampling Type/Medium	Collection Frequency	Total Annual Sample Collections	Analyses Performed/Composite Frequency				
AFFXVRD 3.0 km SSE at Fox Valley	Particulate air samples around the WNYNSC perimeter <u>Reported in:</u> • MTAR • QEMDR • SER	Continuous air particulate filter	→ Weekly	→ 52 each location Weekly filters composited to 4 each location	→ Gross alpha/beta → Quarterly composite for Sr-90, gamma isotopic Total U, U/Pu isotopic, and Am-241 for AFRSPRD and AFGRVAL only				
AFTCORD 3.7 km NNW at Thomas Corners Road									
AFRT240* 2.0 km NE on Route 240									
AFSPRVL 7 km N at Springville									
AFWEVAL 6 km SSE at West Valley									
AFDNKRK** 50 km W at Dunkirk (background)									
AFNASHV** 37 km W at village of Nashville, town of Hanover (background)									
AFBOEHN 2.3 km SW on Dutch Hill Road									
AFRSPRD 1.5 km NW on Rock Springs Road						Continuous desiccant column for water vapor collection	→ Weekly	→ 52 each location (AFRSPRD and AFGRVAL only)	→ H-3
AFGRVAL 29 km S at Great Valley (background)						Continuous charcoal cartridge	→ Monthly	→ 12 composited to 4 each location (AFRSPRD and AFGRVAL only)	→ Quarterly composite for I-129
AFBLKST Bulk Storage Warehouse 2.2 km ESE at Buttermilk Road									

* Filter from duplicate sampler sent to NYSDOH.

** AFNASHV replaced AFDNKRK in 1995.

Sampling Rationale

- AFFXVRD** DOE/EH-0173T, 5.7.4.
AFTCORD
AFRT240 Air samplers put into service by NFS as part of the site's original monitoring program. Perimeter locations chosen to obtain data from places most likely to provide highest concentrations, based on meteorological data.
- AFSPRVL** DOE/EH-0173T, 5.7.4; DOE/EP-0023, 4.2.3.

Off-site (remote) sampler located on private property in nearby community within 15 kilometers of the site (north).
- AFWEVAL** DOE/EH-0173T, 5.7.4; DOE/EP-0023, 4.2.3.

Off-site (remote) sampler located on private property in nearby community within 15 kilometers of the site (southeast).
- AFDNKRK** DOE/EH-0173T, 5.7.4; DOE/EP-0023, 4.2.3.

Off-site (remote) sampler considered to be representative of natural background radiation. Located 50 kilometers west of the site (upwind) on privately owned property. Location discontinued in 1995.
- AFNASHV** DOE/EH-0173T, 5.7.4; DOE/EP-0023, 4.2.3.

Off-site (remote) sampler considered to be representative of natural background radiation. Located 37 kilometers west of the site (upwind) on privately owned property. Location replaced AFDNKRK in 1995.
- AFBOEHN** DOE/EH-0173T, 5.7.4; DOE/EP-0023, 4.2.3.

Perimeter location chosen to obtain data from the place most likely to provide highest elevated release concentrations based on meteorological data. AFBOEHN is located on NYSERDA property at the perimeter.
- AFRSPRD** DOE/EH-0173T, 5.7.4.

Perimeter location chosen to obtain data from the place most likely to provide highest ground-level release concentrations based on meteorological data. AFRSPRD is on WVDP property but outside the main plant operations fence line. I-129 and H-3 are sampled here because the sampling trains were easy to incorporate and the location was most likely to receive effluent releases.
- AFGRVAL** DOE/EH-0173T, 5.7.4; DOE/EP-0023, 4.2.3.

Off-site (remote) sampler considered to be representative of natural background radiation. Located on privately owned property 29 kilometers south of the site (typically upwind). I-129 and H-3 sampled here also.
- AFBLKST** DOE/EH-0173T, 5.7.4.

Off-site monitoring of bulk storage warehouse, near site perimeter.
-

**1995 Monitoring Program
Environmental Surveillance:**

Fallout, Sediment, and Soil

Sample Location Code	Monitoring/Reporting Requirements	Sampling Type/Medium	Collection Frequency	Total Annual Sample Collections	Analyses Performed/Composite Frequency
AFDHFOP 2.3 km SW AFFXFOP 3.0 km SSE AFTCFOP 3.7 km NNW AF24FOP 2.0 km NE ANRGFOP Met tower on-site	Collection of fallout particulate and precipitation around WNYNSC perimeter <u>Reported in:</u> • MTAR • QEMDR • SER	Integrated precipitation	→ Monthly	→ 12 each location	→ Gross alpha/beta, H-3, pH, gamma isotopic
SF Soil Series Surface Soil (at each of ten air samplers)	Long-term fallout accumulation <u>Reported in:</u> • MTAR • QEMDR • SER	Surface plug composite soil	→ Annual	→ 1 each location	→ Gross alpha/beta, gamma isotopic, Sr-90, Pu-239, Am-241, plus U-isotopic and total U at SFRSPRD , SFBOEHN , and SFGRVAL
SFCCSED Cattaraugus Creek at Felton Bridge SFSDSED Cattaraugus Creek at Springville Dam SFBISED Cattaraugus Creek at Bigelow Bridge (background) SFTCSSED Buttermilk Creek at Thomas Corners Road SFBCSED Buttermilk Creek at Fox Valley Road (background)	Deposition in sediment downstream of facility effluents <u>Reported in:</u> • MTAR • QEMDR • SER	Grab stream sediment	→ Annual (Split of SFSDSED and SFBCSED to NYSDOH)	→ 1 each location	→ Gross alpha/beta, gamma isotopic, Sr-90, U/Pu isotopic, total U, Am-241
SN On-site Soil Series: SNSW74A (Near WNSW74A) SNSWAMP (Near WNSWAMP) SNSP006 (Near WNSP006)	<u>Reported in:</u> • MTAR • QEMDR • SER	Surface plug or grab	→ Annual	→ 1 each location	→ Gross alpha/beta, gamma isotopic, Sr-90, Pu-239, Am-241, U-isotopic, total U, Al, Sb, As, Ba, Be, Cd, Ca, Cr, Co, Cu, Fe, Pb, Mg, Mn, Hg, Ni, K, Se, Ag, Na, Tl, V, Zn

Sampling Rationale

AFDHFOP	DOE/EP-0023, 4.7.
AFFXFOP	
AFTCFOP	Collection of fallout particles and precipitation around the site perimeter at established air sampling locations.
AF24FOP	AFDHFOP (Dutch Hill at Boehn road), AFFXFOP (Fox Valley Road), AFTCFOP (Thomas Corners), AF24FOP (Route 240). Indicates short-term effects.
ANRGFOP	Collection of fallout particles and precipitation on-site at the meteorological tower. Indicates short-term effects.
SF Soil Series	DOE/EH-0173T, 5.9.1. Off-site soils collected at air sampling locations.
	SFWEVAL (West Valley), SFFXVRD (Fox Valley Road), SFSPRVL (Springville), SFTCORD (Thomas Corners), SFRT240 (Route 240), SFNASHV (Nashville), SFBOEHN (Boehn Road-Dutch Hill), SFGRVAL (Great Valley), SFRSPRD (Rock Springs Road), SFBLKST (bulk storage warehouse): Collection of long-term fallout data at established air sampler locations via soil sampling.
SFCCSED	DOE/EH-0173T, 5.12.1. Sediment deposition in Cattaraugus Creek at Felton Bridge. Location is first access point of Cattaraugus Creek downstream of the confluence with Buttermilk Creek.
SFSDSED	DOE/EH-0173T, 5.12.1. Sediment deposition in Cattaraugus Creek at Springville Dam. Reservoir provides ideal settling and collection location for sediments downstream of Buttermilk Creek confluence. Located downstream of SFCCSED .
SFBISED	DOE/EH-0173T, 5.12.1. Sediment deposition in Cattaraugus Creek at Bigelow Bridge. Location is upstream of the Buttermilk Creek confluence and serves as a Cattaraugus Creek background location.
SFTCSED	DOE/EH-0173T, 5.12.1. Sediment deposition in Buttermilk Creek at Thomas Corners immediately downstream of all facility liquid effluents.
SFBCESED	DOE/EH-0173T, 5.12.1. Sediment deposition in Buttermilk Creek upstream of facility effluents (background).
SN Soil Series	DOE/EH-0173T, 5.9.1. On-site soil. (Samples may be partially composed of sediments.)
SNSW74A	Surface soil near WNSW74A . Location to be specifically defined by geographic coordinates. Corresponds to site drainage pattern flow (i.e., most likely area of radiological deposition/accumulation).
SNSWAMP	Surface soil near WNSWAMP . Location to be specifically defined by geographic coordinates. Corresponds to site drainage pattern flow (i.e., most likely area of radiological deposition/accumulation).
SNSP006	Surface soil near WNSP006 . Location to be specifically defined by geographic coordinates. Corresponds to site drainage pattern flow (i.e., most likely area of radiological deposition/accumulation).

**1995 Monitoring Program
Environmental Surveillance:**

Off-site Biological

Sample Location Code	Monitoring/Reporting Requirements	Sampling Type/Medium	Collection Frequency	Total Annual Sample Collections	Analyses Performed/ Composite Frequency
BFFCATC Cattaraugus Creek downstream of the Buttermilk Creek confluence	Fish in waters up- and downstream of facility effluents <u>Reported in:</u> • MTAR • QEMDR • SER	Individual collection, biological	Semiannual (Samples at BFFCATC and BFFCTRL shared with NYSDOH)	→ 20 fish each location	→ Gamma isotopic and Sr-90 in edible portions of each individual fish
BFFCTRL Control sample from nearby stream not affected by the WVDP (7 km or more upstream of site effluent point; background)					
BFFCATD Cattaraugus Creek downstream of Springville Dam			Annual (BFFCATD only)	→ 10 fish	→ Gamma isotopic and Sr-90 in edible portions of each individual fish
BFMREED Dairy farm, 3.8 km NNW	Milk from animals foraging around facility perimeter and at background sites <u>Reported in:</u> • MTAR • QEMDR • SER	Grab biological	→ Monthly (BFMREED , BFMCOBO , BFMCTLS , BFMCTLN . Samples at BFMREED and BFMCOBO shared with NYSDOH)	→ 12 monthly samples composited to 4 each location	→ Quarterly composite for gamma isotopic, Sr-90, H-3, and I-129
BFMCOBO Dairy farm, 9 km WNW					
BFMCTLS Control location 25 km S (background)					
BFMCTLN Control location 30 km N (background)					
BFMWIDR Dairy farm, 3.5 km SE of site			Annual (BFMWIDR , BFMSCHT)	→ 1 each location	→ Gamma isotopic, Sr-90, H-3, and I-129
BFMSCHT* Dairy farm 4.8 km S					

* BFMSCHT replaces former location (BFMHUR), which no longer provides milk commercially.

Sampling Rationale

BFFCATC DOE/EH-0173T, 5.11.1.1.

BFFCATD

Radioactivity may enter a food chain in which fish are a major component and are consumed by the local population.

BFFCTRL

Control fish sample to provide background data for comparison with fish caught downstream of facility effluents.

BFMREED DOE/EH-0173T, 5.8.2.1.

BFMCOBO

BFMWIDR Milk from animals foraging around facility perimeter. Milk is consumed by all age groups and is frequently the most important food that could contribute to the radiation dose. Dairy animals pastured near the site and
BFMSCHT at two background locations allow adequate monitoring.

BFMCTLS

Control milk samples collected far from site to provide background data for comparison with near-site milk.

BFMCTLN

**1995 Monitoring Program
Environmental Surveillance:**

Off-site Biological

Sample Location Code	Monitoring/Reporting Requirements	Sampling Type/Medium	Collection Frequency	Total Annual Sample Collections	Analyses Performed/ Composite Frequency
BFVNEAR* Nearby locations BFVCTRL* Remote locations (16 km or more from facility; background) BFHNEAR Beef cattle/milk cow forage from near-site location BFHCTLS or BFHCTLN Beef cattle/milk cow forage from control location south or north (background)	Fruit and vegetables grown near facility perimeter, downwind if possible <u>Reported in:</u> • MTAR • QEMDR • SER	Grab biological (fruits and vegetables) Grab biological	→ Annual, at harvest (BFVNEAR and BFVCTRL) → Annual (BFHNEAR , BFHCTLS , or BFHCTLN)	→ 3 each (split with NYSDOH) → 1 each location	→ Gamma isotopic and Sr-90 analysis of edible portions, H-3 in free moisture → Gamma isotopic, Sr-90
BFBNEAR Beef animal from nearby farm in downwind direction BFBCTRL Beef animal from control location 16 km or more from facility (background)	Meat (beef foraging near facility perimeter, downwind if possible) <u>Reported in:</u> • MTAR • QEMDR • SER	Grab biological	→ Semiannual	→ 2 each location	→ Gamma isotopic and Sr-90 analysis of meat, H-3 in free moisture
BFDNEAR Deer in vicinity of the site BFDCTRL Control deer 16 km or more from facility (background)	Meat (deer foraging near facility perimeter) <u>Reported in:</u> • MTAR • QEMDR • SER	Individual collection biological	→ Annual, during hunting season (BFDNEAR sample split with NYSDOH) During year as available (BFDCTRL sample split with NYSDOH)	→ 3 → 3	→ Gamma isotopic and Sr-90 analysis of meat, H-3 in free moisture → Gamma isotopic and Sr-90 analysis of meat, H-3 in free moisture

* Corn, apple, and bean samples are identified specifically as follows: corn = BFVNEAC and BFVCTRC; apples = BFVNEAA and BFCTRA; beans = BFVNEAB and BFVCTRB.

Sampling Rationale

BFVNEAR DOE/EH-0173T, 5.8.2.2.

Fruits and vegetables (corn, apples, and beans) collected from areas near the site. Collected, if possible, from areas near the site predicted to have worst case downwind concentrations of radionuclides in air and soil. Sample analysis reflects steady state/chronic uptake or contamination of foodstuffs as a result of site activities. Possible pathway to humans or indirectly through animals.

BFVCTRL DOE/EH-0173T, 5.8.2.2.

Fruits and vegetables collected from area remote from the site. Background fruits and vegetables collected for comparison with near-site samples. Collected in area(s) of no possible site impact.

BFHNEAR DOE/EH-0173T, 5.8.2.2.

Hay collected from areas near the site. Same as for near-site fruits and vegetables (**BFVNEAR**). Indirect pathway to humans through animals. Collected with either beef or milk sample location.

BFHCTLS DOE/EH-0173T, 5.8.2.2.

BFHCTLN

Hay collected from areas remote from the site. Background hay collected for comparison with near-site samples. Collected in area(s) of no possible site impact.

BFBNEAR DOE/EH-0173T, 5.8.2.3.

Beef collected from animals raised near the site. Following the rationale for vegetable matter collected near site (**BFVNEAR** and **BFHNEAR**), edible flesh portion of beef animals is analyzed to determine possible radionuclide content passable directly to humans. For animals foraging downwind in areas of maximum probable site impact.

BFBCTRL DOE/EH-0173T, 5.8.2.3.

Beef collected from animals raised far from the site. Background beef collected for comparison with near-site samples. Collected in area(s) of no possible site impact.

BFDNEAR DOE/EH-0173T, 5.8.3.

Venison from deer herd found living near the site. Same as for beef (**BFBNEAR**).

BFDCTRL DOE/EH-0173T, 5.8.3.

Venison from deer herd living far from the site. Background deer meat collected for comparison with near-site samples. Collected in area(s) of no possible site impact.

**1995 Monitoring Program
Environmental Surveillance:**

Off-site Direct Radiation

Sample Location Code	Monitoring/Reporting Requirements	Sampling Type/Medium	Collection Frequency	Total Annual Sample Collections	Analyses Performed/ Composite Frequency
<p>DFTLD Series Thermoluminescent Dosimetry (TLD) Off-site:</p> <p>#1-16 At each of 16 compass sectors, at nearest accessible perimeter point</p> <p>#17 "5 Points" landfill, 19 km SW (background)</p> <p>#20 1,500 m NW (downwind receptor)</p> <p>#21 Springville 6 km N</p> <p>#22 West Valley 6 km SSE</p> <p>#23 Great Valley 29 km S (background)</p> <p>#37 Dunkirk 50 km NW - early 1995 (background)</p> <p>Nashville 37 km NW - late 1995 (background)</p> <p>#41 Sardinia-Savage Road 24 km NE (background)</p>	<p>Direct radiation around facility</p> <p><u>Reported in:</u></p> <ul style="list-style-type: none"> • QEMDR • SER 	<p>Integrating LiF TLD</p>	<p>→ Quarterly</p>	<p>→ 5 TLDs at each of 23 locations collected 4 times per year</p>	<p>→ Quarterly gamma radiation exposure</p>

Sampling Rationale

DOSIMETRY DOE/EH-0173T, 5.5 and DOE/EP-0023, 4.6.3.

Off-site

TLDs offer continuous integrated environmental gamma-ray monitoring and have been deployed systematically about the site. Off-site TLDs are used to verify that site activities have not adversely affected the surrounding environs.

In addition to general NRC crosschecks at selected sites, a biennial HPIC gamma radiation measurement is completed at all TLD locations.

**1995 Monitoring Program
Environmental Surveillance:**

On-site Direct Radiation

Sample Location Code	Monitoring/Reporting Requirements	Sampling Type/Medium	Collection Frequency	Total Annual Sample Collections	Analyses Performed/ Composite Frequency
DNTLD Series Thermoluminescent Dosimetry (TLD) On-site: #18, #19, #33 At three corners of SDA #24, #26-32, #34 (9) At security fence around site #35, #36, #38-40 (5) On-site near operational areas #25 Rock Springs Road 500 m NNW of plant #42 SDA T-1 Building #43 SDA West Perimeter Fence	Direct radiation on facility grounds <u>Reported in:</u> • QEMDR • SER	Integrating LiF TLD	→ Quarterly	→ 5 TLDs at each of 20 sites collected 4 times per year	→ Quarterly gamma radiation exposure

Sampling Rationale

DOSIMETRY DOE/EH-0173T, 5.4 and 5.5.

On-site

On-site TLDs monitor waste management units and verify that the potential dose rate to the general public (i.e., Rock Springs Road) is below 100 mrem/annum (1 mSv/annum) from site activities.

In addition to general NRC crosschecks at selected sites, a biennial HPIC gamma radiation measurement is completed at all locations.

Potential TLD sampling locations are continually evaluated with respect to site activities.




Figure A-1. On-site Air Monitoring and Sampling Points.




Figure A-2. Sampling Locations for On-site Surface Water and Soil.




Figure A-3. On-Site Groundwater Monitoring Network (Includes wells not actively monitored following May 1995 but retained for water level measurements).




Figure A-4. Location of Off-site Surface Water and Sediment Samples.




Figure A-5. Near-site Drinking Water and Biological Sample Points.




Figure A-6. Location of Perimeter Air, Soil, and Fallout Sampling Points.




Figure A-7. Location of Off-site Thermoluminescent Dosimeters (TLDs).




Figure A-8. Location of On-site Thermoluminescent Dosimeters (TLDs).

